







2010

## Recommended Immunizations for Babies

 at birth	HepB	
 2 months	HepB + DTaP + PCV + Hib + Polio + RV 1-2 mos	
 4 months	DTaP + PCV + Hib + Polio + RV	
 6 months	HepB + DTaP + PCV + Hib + Polio + RV 6-18 mos*	(Influenza) 6 mos through 18 years* seasonal** 2009 H1N1***
 12 months	MMR + PCV + Hib + Varicella + HepA 12-15 mos* 12-15 mos* 12-15 mos* 12-15 mos* 12-23 mos*	(Influenza) 6 mos through 18 years* seasonal** 2009 H1N1***
 15 months	DTaP 15-18 mos*	(Influenza) 6 mos through 18 years* seasonal** 2009 H1N1***

## Vaccine Descriptions:

HepB: protects against hepatitis B

DTaP: a combined vaccine that protects against diphtheria, tetanus, and pertussis (whooping cough)

Hib: protects against *Haemophilus influenzae* Type b

PCV: protects against pneumococcal disease

Polio: protects against polio, the vaccine is also known as IPV

RV: protects against infections caused by the Rotavirus

Influenza: protects against influenza (flu)

MMR: protects against measles, mumps, and rubella (German measles)

Varicella: protects against varicella, also known as chickenpox

HepA: protects against hepatitis A

**NOTE:** If your children miss a shot, you don't need to start over, just go back to your doctor for the next shot. The doctor will help you keep your children up-to-date on his or her vaccinations.

\* This is the age range in which this vaccine should be given.


\*\* Influenza is a seasonal vaccine. All children ages 6 months through 18 years should receive vaccination during the influenza season each year. If this is the first time for flu vaccine, a child should receive two doses, separated by at least 4 weeks. If a child only receives one dose in the first season, he or she should receive two doses the next season.

\*\*\* In addition to seasonal influenza vaccine, children also are recommended to receive the 2009 H1N1 influenza vaccine. Children younger than ten years should receive two doses of this vaccine separated by approximately 1 month.

For more information, call toll free 1-800-CDC-INFO (1-800-232-4636) or visit [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines)

The Recommended Immunization Schedules for Persons Aged 0 Through 18 Years are approved by the Advisory Committee on Immunization Practices ([www.cdc.gov/vaccines/recs/acip](http://www.cdc.gov/vaccines/recs/acip)), the American Academy of Pediatrics (<http://www.aap.org>), and the American Academy of Family Physicians (<http://www.aafp.org>).

Parents, don't let your child get left behind!



## School Year 2010-2011

Oregon law requires the following shots for school and child care attendance\*

A child 18 months or older entering  
**Preschool, Child Care, or  
Head Start** needs\*

4 Diphtheria/Tetanus/Pertussis (DTaP)  
3 Polio  
1 Varicella (chickenpox)  
1 Measles/Mumps/Rubella (MMR)  
3 Hepatitis B  
2 Hepatitis A  
3 or 4 Hib

A student entering  
**Kindergarten, 1<sup>st</sup> or 2<sup>nd</sup>  
Grade** needs\*

5 Diphtheria/Tetanus/Pertussis (DTaP)  
4 Polio  
1 Varicella (chickenpox)  
2 Measles  
1 Mumps  
1 Rubella  
3 Hepatitis B  
2 Hepatitis A

A student entering **Grades 3-6  
or Grades 10-12** needs\*

5 Diphtheria/Tetanus/Pertussis (DTaP)  
4 Polio  
1 Varicella (chickenpox)  
2 Measles  
1 Mumps  
1 Rubella  
3 Hepatitis B

A student entering  
**Grades 7-9** needs\*

5 Diphtheria/Tetanus/Pertussis (DTaP)  
1 Tdap  
4 Polio  
1 Varicella (chickenpox)  
2 Measles  
1 Mumps  
1 Rubella  
3 Hepatitis B

\* At all ages and grades, the number of doses required varies by a child's age and how long ago they were vaccinated. Please check with your child's school, childcare or healthcare provider for details. (1/2010)

# Recommended Adult Immunization Schedule UNITED STATES - 2010

Note: These recommendations *must* be read with the footnotes that follow containing number of doses, intervals between doses, and other important information.

**Figure 1. Recommended adult immunization schedule, by vaccine and age group**

VACCINE ▼	AGE GROUP ▶	19–26 years	27–49 years	50–59 years	60–64 years	≥65 years
Tetanus, diphtheria, pertussis (Td/Tdap) <sup>1,*</sup>		Substitute 1-time dose of Tdap for Td booster; then boost with Td every 10 yrs				Td booster every 10 yrs
Human papillomavirus (HPV) <sup>2,*</sup>		3 doses (females)				
Varicella <sup>3,*</sup>		2 doses				
Zoster <sup>4</sup>					1 dose	
Measles, mumps, rubella (MMR) <sup>5,*</sup>		1 or 2 doses		1 dose		
Influenza <sup>6,*</sup>		1 dose annually				
Pneumococcal (polysaccharide) <sup>7,a</sup>		1 or 2 doses				1 dose
Hepatitis A <sup>9,*</sup>		2 doses				
Hepatitis B <sup>10,*</sup>		3 doses				
Meningococcal <sup>11,*</sup>		1 or more doses				

\*Covered by the Vaccine Injury Compensation Program.

For all persons in this category who meet the age requirements and who lack evidence of immunity (e.g., lack documentation of vaccination or have no evidence of prior infection)

Recommended if some other risk factor is present (e.g., on the basis of medical, occupational, lifestyle, or other indications)

No recommendation

Report all clinically significant postvaccination reactions to the Vaccine Adverse Event Reporting System (VAERS). Reporting forms and instructions on filing a VAERS report are available at [www.vaers.hhs.gov](http://www.vaers.hhs.gov) or by telephone, 800-822-7967.

Information on how to file a Vaccine Injury Compensation Program claim is available at [www.hrsa.gov/vaccinecompensation](http://www.hrsa.gov/vaccinecompensation) or by telephone, 800-338-2382. To file a claim for vaccine injury, contact the U.S. Court of Federal Claims, 717 Madison Place, N.W., Washington, D.C. 20005; telephone, 202-357-6400.

Additional information about the vaccines in this schedule, extent of available data, and contraindications for vaccination is also available at [www.cdc.gov/vaccines](http://www.cdc.gov/vaccines) or from the CDC-INFO Contact Center at 800-CDC-INFO (800-232-4636) in English and Spanish, 24 hours a day, 7 days a week.

Use of trade names and commercial sources is for identification only and does not imply endorsement by the U.S. Department of Health and Human Services.